ONE STEP CLOSER CO™

BY KAROLINA NOWICKA

EMERGENCY TOOLKIT · NO. 03

Emotional *Flooding*

What to do when emotion overwhelms your ability to think or function

Too Much · Can't Think · Can't Stop · Overwhelmed

The Nervous System Regulation Toolkit Series

One Step Closer Co™

© Karolina Nowicka. All rights reserved.
No part of this toolkit may be reproduced, distributed,
or transmitted in any form without prior written permission.

This is a trauma-informed self-help and educational resource. It is not therapy, medical advice, diagnosis, treatment, or crisis support. Use only the tools that feel safe and appropriate for your circumstances. Seek qualified professional or emergency support when needed.

ONESTEPCLOSERCO.COM

BEFORE YOU BEGIN

A Note Before You Begin

Emotional flooding is not weakness. It is not drama. It is your nervous system doing exactly what it was built to do when it detects that the emotional load has exceeded what can be held.

During intense emotional activation, reasoning, language, planning, and impulse control may become temporarily harder to access. You are not being irrational. You are being human, in a state where rational thought is temporarily unavailable.

> *Reasoning alone is often not enough during intense flooding. Body-based grounding and reduced stimulation may make reflective thinking more accessible again.*

These tools don't stop you from feeling. They create enough physiological space for your thinking brain to come back online — so the feeling can move through you rather than take you under.

The goal is not calm. The goal is one notch less flooded than you are right now.

`You are not too much. You are overloaded. There is a difference.`

Flooding can also occur during real danger, abuse, intoxication, withdrawal, medical illness, or psychiatric crisis. Do not assume every intense state is merely emotional activation. When safety or health is uncertain, seek appropriate support.

ORIENTATION

How to Use This Toolkit

This one is different. When you are flooded, reading is hard. So this toolkit is built to be used in very small pieces. This toolkit contains six core tools, supported by a three-step protocol, quick-reference guide, personal profile, and customized action plan.

THREE WAYS TO USE IT

> IN THE MOMENT
>
> Go straight to the Quick Reference. Pick the shortest tool. Do just that. Reading comes later.

> LEARNING MODE
>
> Read through once when calm. The understanding of what flooding actually is makes the tools work better.

> BUILDING A PRACTICE
>
> Complete My Flooding Profile and My Personal Protocol. The more you know your pattern, the faster you can interrupt it.

FIND YOUR TOOL BY WHAT'S HAPPENING RIGHT NOW

Need something physical, fast → Tool 01: Cool Sensory Reset	Spinning, can't land anywhere → Tool 02: 5-4-3-2-1 Grounding
Body won't stop, needs to move → Tool 03: Movement Discharge	Breath is short and tight → Tool 04: Breath Regulation
Can't name what this even is → Tool 05: Emotional Labeling	Flooding keeps returning → Tool 06: The Container

Important: when flooded, do not make decisions, send messages, or have important conversations. The tool comes first.

UNDERSTANDING FLOODING

What Is Emotional Flooding?

Daniel Siegel describes a Window of Tolerance — the zone where we can function, feel, and think at the same time. Flooding happens when emotion pushes us above that window.

The Window of Tolerance is a widely used clinical framework for describing the range in which a person can remain emotionally present while still thinking and functioning. It is a useful model, not a precise measurement or diagnosis.

HIGHER ACTIVATION / HYPERAROUSAL
Overwhelmed · Can't think · Rage · Panic · Crying · Spinning

WORKABLE RANGE / WINDOW OF TOLERANCE
Can feel and think at the same time

LOWER ACTIVATION / HYPOAROUSAL (see Toolkit 04)
Shutdown · Numb · Flat · Disconnected · Collapsed

Reasoning, planning, and impulse control may become harder to access during intense activation. This is not a choice. It is a biological response to overwhelm.

SIGNS YOU ARE FLOODED

- Can't finish a sentence
- Crying that won't stop
- Rage with no off switch
- Heart pounding fast
- Mind completely blank
- Everything feels urgent
- Can't hear what's said
- Shaking or trembling

The most important thing to know when flooded: you are not broken and you are not dangerous. You are temporarily above your window. The tools bring you back.

EMERGENCY PROTOCOL

The 3-Step Emergency Protocol

When flooded, decision-making is compromised. This sequence removes the need to decide anything complicated. Follow the steps in order.

01 PAUSE NONESSENTIAL ACTION

Put down the phone when safe to do so. Pause the conversation, scrolling, drafting, or decision-making. Do not stop an activity when doing so would create danger, such as while driving, supervising a child, cooking over an active flame, or handling machinery.

02 COOL THE BODY

Cold water on your face or wrists. If unavailable, press both feet hard into the floor and hold something with firm pressure. The goal is a physical signal that may help redirect attention away from the flood.

03 CHOOSE ONE TOOL

Do not choose the best tool. Choose the first tool that comes to mind from the Quick Reference. One tool for 60 seconds is enough to begin the return.

The single most important rule when flooded: do not make decisions, send messages, or have significant conversations until you are back in your window. What feels urgent almost never is.

→ If the flooding is connected to a safety concern — yours or someone else's — stop here and contact appropriate support immediately.

→ Cold techniques are optional. Avoid extreme cold or sudden immersion. People with cardiovascular conditions, fainting history, circulation problems, pregnancy, or other medical concerns should use a neutral grounding alternative unless cleared by a professional.

TOOL 01

Cool Sensory Reset

01

Physiological interrupt

USE WHEN: Need something physical, fast

A noticeable change in temperature may help redirect your attention to the present moment. This works well for many people, though not everyone finds cold or cool sensations helpful or accessible.

OPTION A — COOL WATER

1. Run cool water over your hands and wrists, or splash it gently on your face.
2. Let the sensation register fully for a few seconds before drying off.
3. Repeat once or twice if it helps.

OPTION B — COOL CLOTH OR OBJECT

1. Press a cool, damp cloth to your face, neck, or wrists.
2. Or hold something cool and solid: a can, a smooth stone, a countertop.
3. Notice the temperature and texture under your hand.

OPTION C — FRESH AIR

1. Step outside or open a window, if that is available to you.
2. Let the air move across your face and hands for a moment.
3. Take a few slow breaths in the cooler air.

→ This works through attention and sensation, not through a guaranteed physiological override. If cold or cool sensations feel unpleasant or are medically inadvisable for you, choose a different tool.

TOOL 02

5-4-3-2-1 Grounding

02

Sensory anchoring to the present

USE WHEN: Spinning, can't land anywhere

When flooded, the mind races between past and future, between catastrophe and meaning. This tool interrupts that by anchoring attention to what is physically real and present right now.

Move slowly through each sense. This is not a race. The slower you go, the more effective it is.

5 THINGS YOU CAN SEE

Look around deliberately. Name five things out loud or in writing. Be specific: not 'a chair' but 'a gray chair with a red cushion.'

4 THINGS YOU CAN TOUCH

Make physical contact with four things. Notice texture, temperature, weight. Press into them slightly.

3 THINGS YOU CAN HEAR

Go still and listen. Name three sounds — near and far. Background noise counts.

2 THINGS YOU CAN SMELL

Breathe in. Name two smells. If nothing registers, find something — coffee, fabric, your own skin.

1 THING YOU CAN TASTE

Notice what is in your mouth. Have a sip of water if you need something concrete.

> *Grounding works because you cannot be fully in the past or future and fully in your senses at the same time. The senses live in the present moment.*

> *You do not have to complete all five categories. Naming one thing you see and one sensation you feel can be enough. If scanning the environment increases overwhelm, choose one neutral object and describe only its shape, color, and texture.*

TOOL 03

Movement Discharge

03

Releasing the flood through the body

USE WHEN: Body won't stop, needs to move

Gentle movement may provide the activation with a safe direction and help reduce the sense of being trapped inside it.

CHOOSE ONE — DO IT FOR 60–120 SECONDS

Shake

Stand and shake your hands, arms, legs. Let the whole body tremble slightly. Animals do this naturally after threat — it completes the stress cycle.

Walk fast

Leave the space you're in. Walk with intention — arms swinging, pace purposeful. Outside is better. Around the block is enough.

Push into a wall

Stand close. Press both palms flat against the wall. Push as hard as you can for 10 seconds. Release. Repeat 3 times. This can give intense energy a safe outlet.

Cold + movement

Splash your face, then walk immediately. The combination of dive reflex and movement is the fastest known return to the window.

Rhythmic tapping

Alternate tapping left knee, right knee, steadily. Add sound if it helps. Bilateral stimulation can help organize a flooded nervous system.

Stomp and hum

Stamp both feet alternately on the floor while humming any note. Sounds playful, and can help organize a flooded nervous system through vagal activation.

→ Every movement here has a smaller version: shake or tap while seated, or simply tense and release instead of standing. Nothing here should be forced. Stop for pain, dizziness, or worsening symptoms.

TOOL 04

Breath Regulation

04

Using the breath to lower the flood

USE WHEN: Breath is short and tight

During flooding, breathing becomes shallow, fast, or held entirely. This amplifies the physiological state of overwhelm. Deliberately changing the breath pattern sends a direct signal to the nervous system that the threat has passed.

Important: do not force deep breathing if you are very flooded — it can increase panic. Start with just slowing the exhale.

PHYSIOLOGICAL SIGH

Double inhale through the nose (sniff, then a second smaller sniff to top up the lungs). Long, slow exhale through the mouth. The double inhale deflates the tiny air sacs that collapse during stress. One to three repetitions is enough.

EXTENDED EXHALE

Inhale for 4 counts. Exhale for 8 counts. Do not force the inhale — the exhale is where the regulation happens. A longer, slower exhale may activate the parasympathetic brake. Repeat 5–8 rounds, or fewer if that feels like enough.

BOX BREATH

Inhale gently. Hold only if that feels comfortable — a shorter hold or no hold at all works too. Exhale slowly. Repeat, adjusting the counts to whatever feels manageable right now.

→ Humming can lengthen the exhale if that feels comfortable — it is optional. No breath holding is required for any of these. If focusing on your breathing increases panic, dizziness, air hunger, or discomfort, stop and use external grounding instead.

TOOL 05

05

Emotional Labeling

Name it to tame it

USE WHEN: Can't name what this even is

Neuroscientist Dan Siegel coined the phrase 'name it to tame it.' Research shows that putting a specific label on an emotional experience reduces activity in the amygdala and increases prefrontal cortex engagement — literally bringing the thinking brain back online.

The more specific the label, the more effective the regulation. 'Bad' is less useful than 'humiliated.' 'Upset' is less useful than 'terrified and then furious.'

THE PRACTICE

1. Ask: "What am I actually feeling right now?" Try naming the broad category first: scared, angry, sad, ashamed, overloaded, disconnected, or unsure. "I do not know yet" is also a valid answer.
2. Push for specificity. Not 'bad' — but what flavor of bad? Scared? Betrayed? Ashamed? Helpless? Furious?
3. Write it down if possible. The act of writing externalizes the emotion and creates a small but real distance.
4. Name all the layers: "I'm angry, and underneath the anger I'm scared, and underneath the fear I'm deeply sad."
5. Say it out loud: "I feel _____." Speaking activates different neural pathways than thinking.
6. Then add: "And that makes sense, because _____." Self-understanding accelerates regulation.

FROM VAGUE TO SPECIFIC

"I feel bad"

→ Ashamed / Humiliated / Guilty / Disappointed

"I feel upset"

→ Terrified / Furious / Devastated / Overwhelmed

"I feel weird"

→ Dissociated / Numb / Unreal / Disconnected

You are not your emotion. You are the one experiencing the emotion. Naming it reminds you of that difference.

TOOL 06

The Container

06

When the flood keeps returning

USE WHEN: Flooding keeps returning

Sometimes the flood isn't one wave — it keeps coming back. Repeated flooding often means the nervous system doesn't yet believe the emotion has anywhere safe to go. The Container gives it a place.

THE IMAGERY PRACTICE

1. Close your eyes or soften your gaze. Take three slow breaths.
2. Imagine a container. Make it solid, strong, and large enough. A vault, a chest, a deep safe room — whatever feels secure to you.
3. In your mind, place the flooding emotion into the container. You are not destroying it. You are giving it somewhere to wait while you regulate.
4. Close the container. You decide what closes it — a lock, a seal, a key. Notice that you hold the key.
5. Say: "I am not leaving this feeling behind. I am setting it down long enough to come back to my window."
6. When you are ready — in an hour, or tomorrow — you can open the container deliberately, with support if needed.
   I will return to this on _____ at _____.

This tool is not suppression. Suppression ignores and pushes away. Containment acknowledges and chooses the timing. The distinction matters: you are not abandoning the feeling. You are pacing your access to it.

WHEN TO OPEN THE CONTAINER

1. When you are back in your window of tolerance
2. With a therapist or trusted support person
3. When you have time and space to feel without needing to function immediately after

→ If the imagery doesn't come easily, try this: physically write the feeling on paper, fold it, and put it in a drawer. The physical act works the same way.

AFTER THE FLOOD

After the Flood

Emotional flooding is physiologically exhausting. After the wave passes, the body needs care — not analysis, not processing, not figuring out what it means. Just care.

RECOVERY RITUALS

1. Rest without guilt. The nervous system has been in overdrive. It needs to recover.
2. Eat or drink something warm if possible. The body depletes real resources during a flood.
3. Do not have the conversation or send the message yet. Wait until tomorrow at the earliest.
4. Avoid alcohol. It feels like regulation — it delays it and typically worsens the next flood.
5. Notice what triggered the flood without judging yourself for being triggered.
6. Journal when ready: "What was the emotion beneath the flood? What did it need?"

> *Flooding is not failure. It is information — about capacity, about what matters, about where your edges are. The work is not to become someone who never floods. It is to become someone who knows how to return.*

Over time, with practice, the window of tolerance widens. Floods become shorter. Return becomes faster. Not because you feel less — but because you've learned to trust that you can come back.

QUICK REFERENCE

When to Use What

TOOL 01

Cool Sensory Reset

When: Need something fast

30–60 sec

30–60 sec

TOOL 02

5-4-3-2-1 Grounding

When: Spinning, can't land

3–5 min

3–5 min

TOOL 03

Movement Discharge

When: Body needs to move

1–2 min

1–2 min

TOOL 04

Breath Regulation

When: Breath short and tight

3–8 min

3–8 min

TOOL 05

Emotional Labeling

When: Can't name what this is

5 min

5 min

TOOL 06

The Container

When: Flood keeps returning

5–10 min

5–10 min

ALWAYS FIRST

Pause nonessential action → Cool the body → Choose one tool

PERSONAL WORK

My Flooding Profile

Complete this after the flood has passed. Use it to recognize your pattern faster next time.

What tends to trigger my flooding

Conflict, overwhelm, bad news, criticism, injustice…

How my body signals flooding is starting

Chest tightens, vision narrows, voice changes, hands shake…

What I do when flooded that I regret

Say things, send messages, withdraw, escalate…

What has helped me return

Cold water, moving, a specific person, a phrase, time alone…

The earliest warning sign — before I am fully flooded

The moment I can still choose. What does that moment look like?

What I need other people to know when I am flooded

PERSONAL WORK

My Personal Protocol

Choose your tools and write them in. Made when calm, for when you're not.

I KNOW I AM FLOODING WHEN…

MY THREE TOOLS — IN ORDER

01 FIRST

The tool I reach for immediately:

02 THEN

If the flood continues, I do:

03 FINALLY

If I am still above my window, I use:

WHAT I WILL NOT DO WHILE FLOODED

e.g. Send that message, make that call, make any decision

MY REMINDER PHRASE

What I say to myself when flooding starts:

A FINAL NOTE

You feel deeply.

That is not the problem.

The capacity to feel everything — the grief, the rage, the love, the overwhelm — is not a malfunction. It is how you are made. The flooding is not evidence that you are broken. It is evidence that you are alive and paying attention.

The work is not to become someone who doesn't flood. It is to become someone who knows the flood will pass — and who has learned, slowly, imperfectly, and with enormous courage, how to move through it.

You came back from every flood that came before this one.

You will come back from this one too.

Emotional Flooding

The Nervous System Regulation Toolkit Series · No. 03

One Step Closer Co™ · Karolina Nowicka

ONESTEPCLOSERCO.COM

Self-help resource. Not a replacement for professional care or crisis support.